

SMITHSONIAN INSTITUTION.
UNITED STATES NATIONAL MUSEUM.

ILLUSTRATIONS AND DESCRIPTIONS OF NEW,
UNFIGURED, OR IMPERFECTLY KNOWN
SHELLS, CHIEFLY AMERICAN, IN
THE U. S. NATIONAL MUSEUM.

BY

WILLIAM HEALEY DALL,

Honorary Curator, Division of Mollusks.

From the Proceedings of the United States National Museum, Vol. XXIV, pages 499-566
(with Plates XXVII-XL).

[No. 1264.]



WASHINGTON:
GOVERNMENT PRINTING OFFICE.

1902.

Color pale yellowish, with darker yellow or brown flammules. Only two imperfect specimens of this fine species are known, both from the above locality.

TEREBRA (SUBULA) FLORIDANA Dall.

Plate XXIX, fig. 9.

Subula floridana DALL, Rep. Blake Gastr., Bull. Mus. Comp. Zool., XVIII, 1889, p. 63.

Terebra (Subula) floridana DALL, Bull. 37, U. S. Nat. Mus., 1889, p. 94.

Key West and Florida Strait, dredged by the U. S. Fish Commission steamer *Albatross* in 45 to 56 fathoms, sand. U.S.N.M., 87222. It is of a pale straw color.

TEREBRA (ACUS) RUSHII Dall.

Plate XXIX, fig. 6.

Terebra (Acus) rushii DALL, Rep. Blake Gastr., Bull. Mus. Comp. Zool., XVIII, 1889, p. 64.

Dredged 5 miles off Cape Florida, in 8 fathoms, by Dr. W. H. Rush, U. S. N. U.S.N.M., 82952.

It is a small, brilliantly white shell, remarkable for its exclusively spiral sculpture.

CONUS STIMPSONI, new species.

Plate XXIX, fig. 7.

Conus stimpsoni DALL, MS., in Coll. U. S. Nat. Mus.

Dredged by the U. S. Fish Commission steamer *Albatross*, in 60 fathoms, off Key West. U.S.N.M., 107371.

Shell of about twelve whorls, as figured, slopes of the spire somewhat concave, turreted, the nuclear whorl rounded, smooth; the following four furnished with a beaded keel at the shoulder; this keel becomes entire on the subsequent whorls; above the shoulder the whorls are slightly concave, the suture appressed, and there are about three faint spiral grooves on the concave surface; last whorl with shallow squarish channels, as figured. The periostracum is thin, pale straw color, finely axially striated. The color of the shell is pinkish white, suffused with salmon pink near the shoulder and on the spire and base, with a very faint, cloudy band of the same about midway between base and shoulder. Altitude of shell, 37 mm.

The species is named in honor of the late Dr. William Stimpson, whose last field work was done in the vicinity of the Florida Strait.

DAPHNELLA EUGRAMMATA, new species.

Plate XXIX, fig. 3.

Dredged at station 2328, off Habana, in 203 fathoms, bottom temperature 78° F., by the U. S. Fish Commission. U.S.N.M., 107448.

- Fig. 11. *Ashmunella rhyssa* Dall, profile; diameter 16.0 mm.; p. 500.
 12. *Zonitoides (Pseudohyalina) pugetensis* Dall, from above; major diameter 1.5 mm.; p. 500.
 13. *Ashmunella pseudodonta* Dall, from below; diameter 14.0 mm.; p. 500.
 14. *Ashmunella rhyssa* Dall, from below; diameter 16.0 mm.; p. 500.
 15. *Ashmunella pseudodonta* Dall, from above; diameter 14.0 mm.; p. 500.

PLATE XXVIII.

- Fig. 1. *Ceres nelsoni* Dall, from above; diameter 27.0 mm.; p. 501.
 2. *Holospira (Haplostemma) hamiltoni* Dall, profile of the basal whorls; p. 501.
 3. *Ceres nelsoni* Dall, basal view; diameter 27.0 mm.; p. 501.
 4. *Ashmunella ashmuni* Dall, basal view; diameter 14.0 mm.; p. 501.
 5. Portion of surface of *Ceres nelsoni*, enlarged to show granular sculpture.
 6. *Ashmunella ashmuni* Dall, from above; diameter 14.0 mm.; p. 501.
 7. *Ashmunella pseudodonta* Dall, profile; diameter 14.0 mm.; p. 500.
 8. *Ceres nelsoni* Dall, profile; diameter 27.0 mm.; p. 501.
 9. *Ashmunella ashmuni* Dall, profile; diameter 14.0 mm.; p. 501.
 10. *Siphonaria lineolata* Orbigny, basal view; longitude 24.0 mm.; p. 501.
 11. *Holospira (Haplostemma) hamiltoni* Dall, profile; altitude 19.0 mm.; p. 501.
 12. *Siphonaria alternata* Say, basal view; longitude 24.5 mm.; p. 501.
 13. *Siphonaria lineolata* Orbigny, profile; longitude 24.0 mm.; p. 501.
 14. *Siphonaria alternata* Say, profile; longitude 24.5 mm.; p. 501.

PLATE XXIX.


- Fig. 1. *Cylichna verrillii* Dall; altitude 7.5 mm.; p. 502.
 2. *Retusa mayoi* Dall; altitude 8.3 mm.; p. 502.
 3. *Daphnella eugrammata* Dall; altitude 9.0 mm.; p. 503.
 4. *Admete microscopica* Dall; altitude 4.3 mm.; p. 504.
 5. *Muricidea philippiana* Dall; altitude 17.4 mm.; p. 504.
 6. *Terebra rushii* Dall; altitude 15.0 mm.; p. 503.
 7. *Conus stimpsoni* Dall; altitude 37.0 mm.; p. 503.
 8. *Terebra texana* Dall; altitude 137.0 mm.; p. 502.
 9. *Terebra floridana* Dall; altitude 70.0 mm.; p. 503.
 10. *Scala scipio* Dall; altitude 16.0 mm.; p. 506.
 11. *Aurinia dubia* Broderip; altitude 69.0 mm.; p. 504.

PLATE XXX.

- Fig. 1. *Muricidea (Pseudoneptunea) multangula* Philippi; altitude 31.0 mm.; p. 505.
 2. *Muricidea ostrearum* Conrad; altitude 28.0 mm.; p. 505.
 3. *Scala (Amæa) mitchelli* Dall; altitude 36.0 mm.; p. 506.
 4. The same, basal view; diameter 14.0 mm.; p. 506.
 5. *Umbraculum (Hyalopatina) rushii* Dall; longitude 9.3 mm.; p. 502.
 6. *Latirus cayoehuesonicus* Sowerby and Melvill; altitude 16.0 mm.; p. 505.
 7. *Scala (Cirsotrema) cochlea* Sowerby; altitude 20.0 mm.; p. 506.
 8. *Scala nitidella* Dall; altitude 13.5 mm.; p. 505.
 9. *Scala (Acrilla) retifera* Dall; altitude 12.5 mm.; p. 506.

PLATE XXXI.

- Fig. 1. *Arca adamsi* Smith, var. *conradiana* Dall; longitude 7.0 mm.; p. 508.
 2. *Arca (Cucullaria) sagrinata* Dall; longitude 6.0 mm.; p. 508.
 3. *Pandora (Kennerleyia) bushiana* Dall; longitude 10.0 mm.; p. 511.
 4. *Pandora (Clidiophora) trilineata* Say; longitude 19.0 mm.; p. 511.
 5. *Meretrix (Transennella) conradina* Dall; longitude 13.5 mm.; p. 509.



EAST AMERICAN GASTROPODS.

FOR EXPLANATION OF PLATE SEE PAGE 563.