

[From the REPORT of the BRITISH ASSOCIATION for the ADVANCEMENT of SCIENCE
for 1863.]

WILLIAM H. DALL
SECTIONAL LIBRARY
DIVISION OF MOLLUSKS

FILE 94

SUPPLEMENTARY REPORT

Dall
1896

ON THE

PRESENT STATE OF OUR KNOWLEDGE

WITH REGARD TO

THE MOLLUSCA OF THE WEST COAST
OF NORTH AMERICA.

BY

PHILIP P. CARPENTER, B.A., Ph.D.

LONDON:

PRINTED BY TAYLOR AND FRANCIS, RED LION COURT, FLEET STREET.

1864.

	Nutt.	Jew.	B. A.	Smiths. Ins.	Ken.	Lord.	Swan.	Cooper.
375. <i>Drillia penicillata</i>	—	—	—	L	—	—	—	—
376. — <i>cancellata</i>	—	—	—	—	P	—	—	—
377. <i>Mangelia levidensis</i>	—	—	—	—	P	—	V	—
378. — <i>tabulata</i>	—	—	—	—	—	—	V	—
379. — <i>interfossa</i>	—	—	—	—	—	—	V	—
380. — <i>crebricostata</i>	—	—	—	—	—	—	V	—
381. — <i>variegata</i>	—	B	—	—	—	—	—	—
381 b. — ? <i>var. nitens</i>	—	B	—	—	—	—	—	—
382. — <i>angulata</i>	—	B	—	—	P	—	—	M
383. <i>Bela fidicula</i>	—	—	P	—	P	V	—	—
384. — <i>excurvata</i>	—	—	—	—	P	—	—	—
385. ? <i>Daphnella aspera</i>	—	—	—	M	—	—	—	—
386. ? — <i>filosa</i>	—	B	—	—	—	—	—	—
387. ? — <i>effusa</i>	—	—	—	—	—	—	V	—
388. <i>Conus Californicus</i>	—	B	C	D	—	—	—	DI
389. <i>Obeliscus</i> ? <i>variegatus</i>	—	—	—	L	—	—	—	D
390. <i>Odostomia nuciformis</i>	—	—	—	—	—	—	V	—
390 b. — ? <i>var. avellana</i>	—	—	—	—	—	—	V	—
391. — <i>satura</i>	—	—	—	—	—	—	V	—
391 b. — ? <i>var. Gouldii</i>	—	—	—	—	—	—	V	—
392. — <i>gravida</i>	—	B	—	—	—	—	—	D
393. — <i>inflata</i>	—	—	—	—	—	—	V	—

375. *Drillia penicillata*, n. s. Like *inermis*, with delicate brownish pencillings.
 376. *Drillia** *cancellata*, ? n. s. Like the young of *incisa*, but nodosely cancellated.
 377. *Mangelia levidensis*, n. s. Stumpy, purplish brown, with rough sculpture.
 378. *Mangelia tabulata*, n. s. Stout, strongly shouldered, coarsely cancellated. Pillar abnormally twisted.
 379. *Mangelia interfossa*, n. s. Like *attenuata*, delicately cancellated.
 380. *Mangelia crebricostata*, n. s. Like *septangularis*, with closely set ribs.
 381. *Mangelia variegata*, n. s. Small, slender, thin, zoned with brown: 9 narrow ribs, and strong spiral striæ.
 381 b. *Mangelia* ?*var. nitens*. Glossy: spiral lines almost obsolete.
 382. *Mangelia angulata*, n. s. Shape of *variegata*, but brown, whirls broad, angular.
 383. *Bela fidicula*, Gld. E.E. Very close to *turricula*, var. 8-10 fm. *Lyall*.
 384. *Bela excurvata*, n. s. Like *Trevelliana*: stumpy, Chrysallid.
 385. ?*Daphnella*† *aspera*, n. s. Elongated, with coarse fenestration.
 386. ?*Daphnella*† *filosa*, n. s. Small, diamond-shaped, but rounded periphery; spirally threaded.
 387. ?*Daphnella*† *effusa*, nom. prov. Thin, extremely drawn-out, sculpture faint.

Family *Conidae*.

388. *Conus Californicus*, Hds. Sulph.=*ravus*, Gld. Chestnut, plain.

Suborder PROBOSCIDIFERA. Family *Pyramidellide*.

389. *Obeliscus* ?*variegatus*, n. s. From Gulf fauna. Periphery with spiral groove. Colour-pattern clouded.
 390. *Odostomia nuciformis*, n. s. Very large, solid, Tornatelloid.
 390 b. *Odostomia* ?*var. avellana*. Shape of *conoidalis*.
 391. *Odostomia satura*, n. s. Large, with swollen whirls like *Bithinia similis*.
 391 b. *Odostomia* ?*var. Gouldii*. Taller, base gently rounded.
 392. *Odostomia gravida*, Gld. Otia. Like *conoidalis*, but nucleus minute.
 393. *Odostomia inflata*, n. s. Like large *dolioliformis*: with most minute spiral striulation. Farallone Is. On *Hal. rufescens*, teste Darbshire.

* A peculiar group of species, resembling *Clionella* (marine, teste Stimpson.)

† Generic position of all these doubtful: perhaps they belong to genera not yet eliminated: *filosa* resembling the Eocene forms between *Conus* and *Pleurotoma*.

|| *Clathurella bella*. Hds. Coop!

" *constricta*. Gabb.

" *crystallina* "

* *Mangelia hexagona* Gabb.

† *Daphnella clathrata* "